

Norton Neuroscience Institute Memory Center

Anti-Amyloid Infusions

Where are the infusion centers?

- **Downtown Louisville**
 - 676 S Floyd St., Louisville, KY 40202
 - Phone: (502)629-5153
- **Northeastern Louisville (Brownsboro)**
 - 4955 Norton Healthcare Blvd., Louisville, KY 40241
 - Phone: (502)394-6315
- **Harrison County Hospital (Corydon, IN)**
 - 1263 Hospital Drive, Suite 110, Corydon, IN 47112
 - Phone: (812)734-0912
- **King's Daughter Hospital (Madison, IN)**
 - 1373 North State Rd. 62, Madison, IN 47250
 - Phone: (812)801-0922

How do I prepare for my infusions?

- A driver is required for each infusion.
- Please drink plenty of water to help with vein access.
- Take your usual medications as prescribed.
- No fasting is required.

What are some things I need to know while taking this drug?

- Tell all of your health care providers that you are receiving anti-amyloid therapy. This includes your doctors, nurses, pharmacists, and dentists.
- Tell your NNI Provider if you have any changes in health conditions or medications while on anti-amyloid therapy.

Norton Neuroscience Institute Memory Center

Norton Medical Plaza III – Brownsboro, Suite 301

4915 Norton Healthcare Blvd.

Louisville, KY 40241

(502) 394-6460 Phone

(502) 394-6465 Fax



**NORTON
NEUROSCIENCE
INSTITUTE**

Norton Neuroscience Institute Memory Center

What are some things I need to know while on Anti-Amyloid therapy?

Monitor for infusion reactions. About 26% of patients receiving Lecanemab and 9% of patients receiving Donanemab note symptoms of an infusion reaction. This is most common with the first few infusions, and in some cases can be delayed for up to several hours. Fortunately, these symptoms are typically mild, resolve within 24 hours and can be managed easily with over the counter medications such as acetaminophen (Tylenol) 650mg – 1000mg every 6 hours, or diphenhydramine (Benadryl) 25mg- 50mg every 6 hours.

Mild infusion reactions include: fever; flu-like signs like chills, aches, shaking, or joint pain; upset stomach.

Moderate to severe infusion reactions include: signs of high or low blood pressure like dizziness or passing out, headache, or change in eyesight; abnormal or pounding heartbeat; or trouble breathing.

Tell your healthcare team right away if you have any of these signs or other effects during OR after the infusion.

When to call your provider:

- **If you are experiencing infusion or drug related reactions:**
fever; flu-like symptoms like chills, aches, shaking, or joint pain; upset stomach or throwing up; signs of high or low blood pressure like dizziness or passing out, headache, or change in eyesight; abnormal or pounding heartbeat; or trouble breathing.
- **Inform your healthcare provider:**
 - **If there are any changes to your health conditions or medication,**
 - **If you need to miss an infusion of the drug, or**
 - **If you have a change in care partner.**

****Call 911 or go to your nearest emergency room for any urgent medical needs***

After Hours On Call Provider:

502-394-6460

Norton Neuroscience Institute Memory Center

Norton Medical Plaza III – Brownsboro, Suite 301

4915 Norton Healthcare Blvd.

Louisville, KY 40241

(502) 394-6460 Phone

(502) 394-6465 Fax



**NORTON
NEUROSCIENCE
INSTITUTE**